



Notes on Early Land Plants Today. 46. Transfer of *Vitalianthus urubuensis* (Marchantiophyta, Lejeuneaceae) to *Cheilolejeunea*

YU-MEI WEI^{1,2}, QIONG HE³, S. ROBERT GRADSTEIN⁴, LAURA V. CAMPOS⁵ & RUI-LIANG ZHU^{1,6}

¹Department of Biology, School of Life Science, East China Normal University, 3663 Zhong Shan North Road, Shanghai 200062, China; rlzhu@bio.ecnu.edu.cn

²Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and the Chinese Academy of Sciences, Guilin, Guangxi 541006, China

³Ningbo Binhai International Cooperative School, Ningbo Binhai Education Group Ltd., 11 Mingyue Road, Binhai New City, Beilun, Ningbo, Zhejiang 315830, China

⁴Muséum National d'Histoire Naturelle, Dept. Systématique et Evolution, Case Postale 39, 57 rue Cuvier, 75231 Paris cedex 05, France

⁵Instituto de Ciencias Naturales, Facultad de Ciencias, Universidad Nacional de Colombia, Bogotá, Colombia

⁶Author for correspondence

Vitalianthus was established by Schuster & Giancotti (1993) based on *Drepanolejeunea bischleriana* K.C.Pôrto & Grolle (1987: 301), which is endemic to the Atlantic coast of north-eastern Brazil. This genus is related to *Drepanolejeunea* (Spruce 1884: 76) Schiffner (1893: 119) (Gradstein *et al.* 2001, Gradstein & Costa 2003, He *et al.* 2012a, b) and characterized by the 1) hyaline papilla at the proximal side of lobule tooth, 2) unbroken basal vitta of 3–7 ocelli in leaf lobes, 3) bilobed underleaves with upright lobes, 4) gynoeical innovations with a pycnolejeuneoid leaf sequence, 5) perianth keels unarmed, and 6) absence of specialized cladia (He *et al.* 2012a). In addition to the type species, this genus currently also contains *Vitalianthus urubuensis* Zartman & Ackerman (2002: 267) from the Amazonian rainforests of Brazil and Colombia (Zartman & Ackerman 2002, Mota de Oliveira & ter Steege 2013, Campos *et al.* in prep.), and *V. guangxianus* Zhu *et al.* (2012a: 32). Our examination of type material and recent collections of *Vitalianthus urubuensis* from Brazil and Colombia reveals that the hyaline papilla of the leaf lobule is distal as in *Cheilolejeunea* (Spruce 1884: 79) Schiffner (1893: 118) (Fig. 1E,F). The moniliate, unbroken vitta of ocelli in the leaf lobes, one of the most important features of *Vitalianthus*, also occurs in some species of *Cheilolejeunea*, e.g. in *Cheilolejeunea falsinervis* (Sande Lacoste 1854: 321) Schuster & Kachroo (1961: 56) from Asia and Australasia. It thus appears that *V. urubuensis* is morphologically more similar to *Cheilolejeunea* than to *Vitalianthus*. The morphological affinity between *Vitalianthus urubuensis* and *Cheilolejeunea* is confirmed by our unpublished molecular data (ITS, *trnG* and *trnL-F*; Wei *et al.*, in prep.) which show that *V. urubuensis* and several species of *Cheilolejeunea* form a monophyletic lineage. Therefore, the following new combination is necessary.

Formal treatment

The format of this note follows Söderström *et al.* (2012).

Cheilolejeunea urubuensis (Zartman & I.L.Ackerman) R.L.Zhu & Y.M.Wei, *comb. nov.* (Fig 1)

Basionym:—*Vitalianthus urubuensis* Zartman & I.L.Ackerman, *Bryologist* 105: 267, 2002 (Zartman & Ackerman 2002).

Type:—BRAZIL. Amazonas State. EMBRAPA research station at kilometer 54 of the BR-174 highway (02°30'40" S, 60°01'46" W), elev. ca 150 m; lowland, seasonally wet terra-firma rainforest in capoeira on the bark of *Vismia cayennensis* (Jacq.) Pers., August 2001, Zartman 2626 (holotype INPA; isotype DUKE!).




FIGURE 1. *Cheilolejeunea urubuensis* (Zartman & I.L.Ackerman) R.L.Zhu & Y.M.Wei. A. Portion of plant with perianth. B. Portion of plant, showing underleaves. C. Portion of plant, ventral view. D. Leaf, ventral view. E, F. Leaf lobule, showing a hyaline papilla (arrow). A from *Laura V. Campos 713* (HSNU); B, F from *Laura V. Campos 712* (HSNU); C-E from *C.E. Zartman 2626* (isotype, DUKE).

Note: Within tropical America, *Cheilolejeunea urubuensis* is the only vittate species in the genus..

Acknowledgements

This research was sponsored by the National Natural Science Foundation of China (nos. 31170190, 31370238), the Special Program for the National Basic Work of the Ministry of Science and Technology, China (No. 2012FY110600), and 211 Project for the East China Normal University. We also thank the curator of DUKE for lending the type specimens. Fieldwork in the Colombian Amazon region was supported by COLCIENCIAS, Colombia.

References

- Gradstein, S.R. & Costa, D.P. (2003) The Hepaticae and Anthocerotae of Brazil. *Memoirs of the New York Botanical Garden* 87: 1–318.
- Gradstein, S.R., Churchill S.P. & Salazar-Allen, N. (2001) Guide to the bryophytes of tropical America. *Memoirs of the New York Botanical Garden* 86: 1–577.
[http://dx.doi.org/10.1016/S1290-0796\(02\)85017-9](http://dx.doi.org/10.1016/S1290-0796(02)85017-9)
- He, Q., Wei, Y.-M. & Zhu, R.-L. (2012a) *Vitalianthus guangxianus* (Lejeuneaceae), a new species from Guangxi, China. *Journal of Bryology* 34: 32–36.

- <http://dx.doi.org/10.1179/1743282011y.0000000038>
- He, Q., Zhu, R.-L., Chantanaorrapint, S., Kornochalert, S. & Printarakul, N. (2012b) *Drepanolejeunea laciniata* (Lejeuneaceae), a new species from northern Thailand. *Cryptogamie Bryologie* 33: 291–298.
<http://dx.doi.org/10.7872/cryb.v33.iss3.2012.291>
- Kachroo, O. & Schuster, R.M. (1961) The genus *Pycnolejeunea* and its affinities to *Cheilolejeunea*, *Euosmolejeunea*, *Nipponolejeunea*, *Tuyamaella*, *Siphonolejeunea* and *Strepsilejeunea*. *Journal of the Linnean Society. Botany* 56: 475–511.
<http://dx.doi.org/10.1111/j.1095-8339.1961.tb02542.x>
- Mota de Oliveira, S. & ter Steege, H. (2013) Floristic overview of the epiphytic bryophytes of *terra firme* forests across the Amazon basin. *Acta Botanica Brasilica* 27: 347–363.
<http://dx.doi.org/10.1590/s0102-33062013000200010>
- Pôrto, K.C. & Grolle, R. (1987) *Drepanolejeunea bischleriana* sp. nov. du Brésil. *Cryptogamie: Bryologie, Lichénologie* 8: 301–304.
- Sande Lacoste, C.M. van der (1854) Novae species Hepaticarum ex insula Java detexit Dr. F. Junghuhn. *Nederlandsch Kruidkundig Archief. Verslagen en Mededelingen der Nederlandsche Botanische Vereeniging* 3: 414–424.
- Schiffner, F. (1893) Hepaticae (Lebermoose). In: Engler, A. & Prantl, K. (Ed), *Die Natürlichen Pflanzenfamilien* 1: 1–141. Leipzig.
- Schuster, R.M. & Giancotti, C. (1993) On *Vitalianthus* Schust. & Giancotti, a new genus of Lejeuneaceae. *Nova Hedwigia* 57: 445–456.
- Söderström, L., Hagborg, A. & von Konrat, M. (2012) Notes on Early Land Plants Today. *Phytotaxa* 65: 41–42.
<http://dx.doi.org/10.11646/phytotaxa.112.1.3>
- Spruce, R. (1884) Hepaticae of the Amazon and the Andes of Peru and Ecuador. *Transactions and Proceedings of the Botanical Society. Edinburgh* 15: 1–308.
- Zartman, C.E. & Ackerman, I.L. (2002) A new species of *Vitalianthus* (Lejeuneaceae, Hepaticae) from the Brazilian Amazon. *The Bryologist* 105: 267–269.
[http://dx.doi.org/10.1639/0007-2745\(2002\)105\[0267:ansovl\]2.0.co;2](http://dx.doi.org/10.1639/0007-2745(2002)105[0267:ansovl]2.0.co;2)