

# Linking Improvement of Soil Structure to Soil Carbon Storage Following Invasion by a C<sub>4</sub> Plant *Spartina alterniflora*

Yanghui He,<sup>1,2,3</sup> Xuhui Zhou,<sup>1,4\*</sup> Weisong Cheng,<sup>1</sup> Lingyan Zhou,<sup>1</sup> Guodong Zhang,<sup>2</sup> Guiyao Zhou,<sup>1</sup> Ruiqiang Liu,<sup>1</sup> Junjiong Shao,<sup>1</sup> Kai Zhu,<sup>3</sup> and Weixin Cheng<sup>3</sup>

<sup>1</sup>Center for Global Change and Ecological Forecasting, Shanghai Key Lab for Urban Ecological Processes and Eco-Restoration, School of Ecological and Environmental Sciences, East China Normal University, Shanghai 200241, China; <sup>2</sup>Coastal Ecosystems Research Station of Yangtze River Estuary, Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, The Institute of Biodiversity Science, School of Life Sciences, Fudan University, 2005 Songhu Road, Shanghai 200433, China; <sup>3</sup>Environmental Studies Department, University of California, Santa Cruz, California 95064, USA; <sup>4</sup>Shanghai Institute of Pollution Control and Ecological Security, 1515 North Zhongshan Rd, Shanghai 200437, China

## ABSTRACT

Coastal wetlands are increasingly recognized as important ecosystems for long-term carbon (C) storage. However, how soil aggregation mediates C accumulation and sequestration in these ecosystems remains unclear. Using the <sup>13</sup>C isotope tracer from the invasion of a C<sub>4</sub> plant, *Spartina alterniflora*, into the native ecosystem originally covered by C<sub>3</sub> plants across Eastern Chinese coastal wetlands, we investigated a potential C stabilization process via soil structural protection. We quantified changes in soil aggregates, soil organic carbon (SOC), soil total nitrogen (STN), and natural <sup>13</sup>C isotope abundance within aggregate fractions across a chronosequence of 0-, 4-, 8-, and 12-year *S. alterniflora* invasion. Our results showed that soil aggregate stability increased

significantly along the chronosequence. Meanwhile, SOC and STN concentrations increased with invasion time in the whole soil and aggregate fractions, which were linked to increasing soil aggregate stability. The contribution of *S. alterniflora*-derived SOC increased from 18.96 to 40.24% in the 0–20 cm layer and from 4.66 to 32.04% in the 20–40 cm layer across the chronosequence from 4 to 12 years with the highest proportion observed in macro-aggregates. Our results indicate that invasion of *S. alterniflora* to coastal wetlands can sequester more C largely due to formation and stabilization of soil aggregates by soil structural protection.

**Key words:** coastal wetland; *Spartina alterniflora*; soil aggregates; soil organic carbon; soil total nitrogen; stable carbon isotope.

Received 24 July 2018; accepted 17 September 2018;  
published online 15 October 2018

**Electronic supplementary material:** The online version of this article (<https://doi.org/10.1007/s10021-018-0308-3>) contains supplementary material, which is available to authorized users.

**Author Contributions** YH, XZ, LZ, and JS designed this experimental study. YH, WC, RL, and GZ performed the research. YH analyzed the data. YH, XZ, KZ, and WC wrote the manuscript, and all authors substantially contributed to the revision.

\*Corresponding author; e-mail: xhzhou@des.ecnu.edu.cn

## INTRODUCTION

The increasing atmospheric concentrations of greenhouse gases (GHGs) from human activities are the main drivers of observed climate change

(IPCC 2013). Strategies to mitigate GHG emissions have focused on the conservation of terrestrial carbon (C) sinks (Agrawal and others 2011). Coastal vegetated ecosystems rank among the most efficient C sinks in the biosphere, and conserving and restoring “blue carbon” sinks has been viewed as a promising strategy (Duarte and others 2005). The overall average rate of C sequestration was  $210 \text{ g CO}_2 \text{ m}^{-2} \text{ y}^{-1}$  in coastal wetlands, which is an order of magnitude greater than peatlands (Roulet 2000; Chmura and others 2003). Moreover, coastal wetlands are not a significant source for atmospheric methane ( $\text{CH}_4$ ) compared with non-tidal wetlands, due to the presence of abundant sulfate ( $\text{SO}_4^{2-}$ ) which inhibits  $\text{CH}_4$  production (Choi and Wang 2004). Due to the higher rate of C sequestration, lower  $\text{CH}_4$  emissions, and other ecosystem services they provide (for example, buffering the impacts of sea-level rise and dissipating wave energy), coastal wetlands are among the most valuable ecosystems for mitigating climate change (Mcleod and others 2011; Kirwan and Megonigal 2013).

Given the importance of coastal wetlands to atmospheric C budget, understanding the processes of C accumulation and mechanisms underlying soil organic carbon (SOC) stabilization provides a basis for coastal wetland management and conservation. However, substantial uncertainties exist in the quantification of SOC sources in coastal wetlands due to complicated environment (Harvey and Mannino 2001). SOC in coastal wetlands comes from both external and internal sources, including relict organic materials such as allochthonous particulate matter and dissolved organic matter, and newly produced organic materials from the primary production of local plants (Neubauer 2008; Kennedy and others 2010; Mcleod and others 2011). Recent works have demonstrated the importance of the burial organic C from plant communities in coastal wetlands (Chmura and others 2003; Donato and others 2011; Kelleway and others 2016). For example, an estimate of nearly 50% of C fixed in marine sediments is thought to be from the internal origin at the global level (Duarte and others 2005). Bull and others (1999) reported that the *Spartina anglica* biomass contributed 37–100% to salt marsh sediments at Wytch Farm, Dorset, UK, as a result of local organic matter burial into the sediments with low oxygen conditions (Alongi 2002; Kristensen and others 2008; Fourqurean and others 2012). However, other studies suggested that external C input was the primary source of the C accumulation in coastal wetlands (Kennedy and others 2010), which was ascribed to marsh accretion and

lateral detritus transported by strong and frequent surface water movements (Hemminga and others 1996; Neubauer 2008). These controversial results may limit our ability to predict future C sequestration and stabilization in coastal ecosystems. Therefore, it is necessary to probe the potential mechanisms underlying the SOC accumulation in coastal wetlands.

Soil C storage and turnover are largely determined by soil structure in upland systems (Golchin and others 1997; Wilson and others 2009). SOC is incorporated into the soil matrix in different sizes and shapes through flocculation, cementation, and arrangement of soil particles (Duiker and others 2003). The position of SOC within different physical fractions causes differential accessibility to microbial decomposers and leads to a wide range of SOC pools that vary in stability and dynamics (Six and others 2000a; Liao and others 2006). For example, the macro-aggregates are formed around the fresh organic matter, which serve as a C source for microbial activity and the production of microbial-derived binding agents (Jastrow 1996; Six and others 1998, 2000b). However, the micro-aggregates, which are bound together by roots, fungal hyphae, or polysaccharides generated by the decomposition of organic inputs, contained a greater proportion of older organic matters that is more resistant to decay (Six and Jastrow 2002). Therefore, the dynamics of aggregate formation are closely linked with the stabilization of soil organic matter (Salome and others 2010). Although numerous studies in upland ecosystems have shown that soil aggregation provides physical protection of SOC against rapid decomposition and plays a critical role in SOC turnover (Christensen 2001; Six and others 2004; Abiven and others 2009), the importance of soil structure controlling the SOC sequestration in coastal wetlands remains largely unknown.

*Spartina alterniflora* was introduced to Chinese coastal wetlands for the purpose of ecological engineering, in 1979 from the USA (Wang and others 2006; An and others 2007). The  $\text{C}_4$  plant *S. alterniflora* has spread rapidly and becomes the dominant vegetation type since it was transplanted to the Dongtan wetlands, which were previously dominated by  $\text{C}_3$  vegetation such as *Scirpus mariqueter* and *Phragmites australis* (Chen and others 2004). This shift in vegetation structure provides a unique opportunity to examine the contribution of coastal vegetation to SOC accumulation and to seek the mechanism of SOC stabilization mediated by soil aggregation in coastal wetlands, although salinity, hydrology and oxygen availability may

influence soil C accumulation (Chambers and others 2016; Graves and others 2016). Furthermore, the chronosequence created by the *S. alterniflora* invasion provides a temporal dimension for exploring the mechanisms contributing to the accumulation and stabilization of SOC. In this study, we hypothesized: (1) the *S. alterniflora* invasion would increase soil aggregate stability and soil C and N contents in the whole soil and aggregate fractions and (2) the increasing SOC was mainly derived from inputs of *S. alterniflora*. The specific objectives of this study were to: (1) evaluate the influence of *S. alterniflora* invasion on soil aggregate stability and C and N pools in aggregate fractions and (2) partition C contributions from *S. alterniflora* in the whole soil and aggregate fractions.

## MATERIALS AND METHODS

### Study Area

This study was conducted at Dongtan wetland of Chongming Island, the largest wetland in the Yangtze River estuary, China (31°03′–31°38′ N, 121°47′–122°05′ E, Figure 1). The local climate is a subtropical monsoon, characterized by a mean annual temperature of 15.2°C and a total annual precipitation of 1025 mm. Dongtan wetland is expanding eastward to the sea at a rate of more than 150 m per year due to the continuous silt sedimentation from the Yangtze River (Zhao and others 2009). The dominant vegetation species were the C<sub>3</sub> plants *Scirpus mariqueter* and *Phragmites australis* before the 1980s. The invasive C<sub>4</sub> plant *Spartina alterniflora*, which was intentionally introduced from the USA to Chinese coastal wetlands in 1979 (Qin and Zhong 1992), was transplanted into Dongtan in May 2001 (Chen and others 2004). Since then, it has been rapidly spreading along the Dongtan wetlands.

### Soil and Plant Sampling

Soil and plant samples were collected from *S. alterniflora* communities in June 2013. The invasion time of *S. alterniflora* within the sample region was identified based on Thematic Mapper (TM) satellite images (Wang and others 2015) and historical records. The chronosequence consisted of *S. alterniflora* invaded in 2001 (SA-12), 2005 (SA-8), 2009 (SA-4), and the bare flat (BF, Figure 1). The sampled region was reasonably flat with a less than 0.5 degree slope. Tidal fluctuation at this tested area is regular and semidiurnal.

Soil samples were collected at four sites (BF, SA-4, SA-8, and SA-12) along the chronosequence

(Figure 1). At each site, three sampling plots (3 × 3 m) were randomly established, and three soil samples were taken at each plot by using a 3-cm-diameter stainless steel sampler from the 0–20 and 20–40 cm layers, respectively. Soil samples within the same layer from each plot were mixed thoroughly. Meanwhile, three replicates of plant samples, including leaves, litters, and roots, were collected at each site. All soil and plant samples were filled in sample sack and kept on ice in coolers, immediately transported to the laboratory at Fudan University, and stored at 4°C for further analysis.

### Laboratory Analyses

Every soil sample was physically cleaned of plant and organic debris with tweezers carefully and divided into two subsamples. One subsample was air-dried for 20 days to analyze for aggregate fractions. Another subsample was dried at 55°C for measuring soil organic carbon (SOC), soil total nitrogen (STN), and stable carbon isotopic composition ( $\delta^{13}\text{C}$ ) of the SOC. The plant samples were dried at 65°C, ground by a planetary ball mill, and passed through a 100-mesh sieve for carbon isotopic analysis by using a MAT 253 isotope ratio mass spectrometer (Thermo Electron, Bremen, Germany).


Soil samples were treated with 1 N HCl for 24 h at room temperature to remove the inorganic carbon before further measurements (Lin and others 1999). The  $\delta^{13}\text{C}$  value of each HCl-treated soil and plant samples was measured on a MAT 253 isotope ratio mass spectrometer (Thermo Electron, Bremen, Germany). Precision for  $\delta^{13}\text{C}$  was  $\pm 0.10\text{‰}$  according to repeated measurements of a laboratory working standard (Shenyang Institute of Applied Ecology, Chinese Academy of Sciences). The  $\delta^{13}\text{C}$  of the organic matter is calculated from:

$$\delta^{13}\text{C} = [R_{\text{sample}}/R_{\text{PDB}} - 1] \times 1000 \quad (1)$$

where  $R_{\text{sample}}$  and  $R_{\text{PDB}}$  are the  $^{13}\text{C}/^{12}\text{C}$  ratio of the sample and the Pee Dee Belemnite (PDB) standard (Desjardins and others 1996; Wooller and others 2003), respectively.

### Soil Physical Fractionation

The size distribution of water-stable aggregates (WSAs) and the stability of moist soil were determined using fast wetting (NVFW) techniques under ambient atmospheric pressure (Kemper and Rose- nau 1986; Sun and others 2014). A 100-g soil sample was placed on the top of a nest of five sieves



**Figure 1.** Location of the sampling sites at Dongtan wetland, Chongming Island, Yangtze River estuary, China. SA-4, SA-8, and SA-12 represent *S. alterniflora* community invaded in 2009, 2005, and 2001, respectively; BF represents bare flat.

with 2, 1, 0.5, 0.25, and 0.106 mm openings and submerged in tap water for 2 min at room temperature. The wet sieve instruments (Model: DIK-2012, Daiki Rika Kogyo Co. Ltd., Japan) were then set in oscillating motion 3 cm up and down at a frequency of 0.5 cycle  $s^{-1}$  for 10 min. After that, the resistant fractions ( $> 2$  mm aggregates were not found) on the top of each sieve and the unstable ( $< 0.106$  mm) aggregates were transferred into beakers. Each aggregate size fraction was dried at  $65^{\circ}\text{C}$  to constant weight. The mean weight diameter (MWD, mm) of soil aggregates is calculated as follows:

$$\text{MWD} = \frac{\sum_{i=1}^5 M_i \times D_i}{\sum_{i=1}^5 M_i} \quad (2)$$

where within each of the five classes of diameter ( $i = 1-5$ ),  $D_i$  [mm] is the central diameter of aggregates in each size class (approached as mean size of upper and lower limits of sieve size used: 1.5,

0.75, 0.375, 0.178, and 0.053 mm, respectively, Barto and others 2010);  $M_i$  [g] is the mass of stable soil aggregates isolated within the  $i$ th diameter class. The fractions above 1 mm were corrected by removing the mass of marly platelets.

### Calculation of Carbon Inputs

The proportion  $f$  (%) of *S. alterniflora*-derived C in the soil fractions was calculated using a two-source mixing model (Chiang and others 2004; Cheng and others 2006) as follows:

$$f = (\delta^{13}\text{C}_{\text{new}} - \delta^{13}\text{C}_{\text{old}}) / (\delta^{13}\text{C}_{\text{sa}} - \delta^{13}\text{C}_{\text{old}}) \times 100 \quad (3)$$

where  $\delta^{13}\text{C}_{\text{sa}}$  is the mean  $\delta^{13}\text{C}$  of *S. alterniflora* plant materials, including litters, rhizomes, and roots;  $\delta^{13}\text{C}_{\text{new}}$  is the mean  $\delta^{13}\text{C}$  of SOC in the soil fractions after the invasion of *S. alterniflora*; and  $\delta^{13}\text{C}_{\text{old}}$  is the mean  $\delta^{13}\text{C}$  of SOC in the soil fractions before

the invasion of *S. alterniflora*; here, it represents the mean  $\delta^{13}\text{C}$  of SOC in the bare tidal mudflat soil fractions.

The accumulation rate ( $R$ ,  $\text{g C kg}^{-1} \text{ soil y}^{-1}$ ) of SOC or *S. alterniflora*-derived C in soil physical fractions is calculated as:

$$R = C_{\text{in}}/t \quad (4)$$

where  $C_{\text{in}}$  is assumed to be the increased storage of SOC or *S. alterniflora*-derived C in soil fractions, and  $t$  is the invasion time.

## Statistical Analyses

One-way analysis of variance (ANOVA) was performed to determine differences in soil physical and chemical characteristics in the whole soil and aggregate fractions among each site. Significant differences among different sites were assessed by Fisher's protected least significant difference (LSD) test at  $P = 0.05$  for all the variables. Regression analyses were used to determine the relationships of MWD, SOC, STN, accumulative rates of SOC and *S. alterniflora*-derived C against plant invasion time. The ANOVA and regression analysis were carried out in SPSS Statistics 23 (SPSS, Inc., Chicago, IL, USA).

## RESULTS

### Soil Physicochemical Properties and Soil Aggregate Stability

Along the chronosequence of *Spartina alterniflora* invasion from bare flat to 12-year invasion, soil pH in the *S. alterniflora* community ranged from 8.11 to 8.49 but with no significant difference. However, soil electrical conductivity and salinity in the *S. alterniflora* community were higher than those in bare flat (Table 1). The  $\delta^{13}\text{C}$  values of *S. alterniflora* ranged from  $-13.45$  to  $-14.44\%$  with a mean value of  $-13.86\% \pm 0.10$  (mean  $\pm$  SE, Table S1), showing no significant difference among leaves, litter, and roots. The  $\delta^{13}\text{C}$  values of SOC in the whole soil and aggregate fractions increased along the chronosequence of *S. alterniflora* invasion from bare flat to 12 years in both the surface (0–20 cm) and deep soil layers (20–40 cm). Meanwhile, within aggregate fractions, the  $\delta^{13}\text{C}$  values increased with the particle size at each site, but not at the bare flat (Table 2).

The distribution of soil aggregate fractions as a proportion of the whole-soil mass varied significantly from bare flat to the invasion of *S. alterniflora* with 4–12 years (Figure 2). The mass proportion of

**Table 1.** Soil Physicochemical Characteristics of *Spartina alterniflora* Communities in Dongtan Wetlands, Shanghai, China

Site	pH	EC (ms/cm)	Salinity (PPT)
Bare flat	$8.49 \pm 0.08^a$	$1.96 \pm 0.14^a$	$5.96 \pm 0.46^a$
SA-4	$8.11 \pm 0.15^a$	$2.76 \pm 0.03^{bc}$	$8.55 \pm 0.09^b$
SA-8	$8.38 \pm 0.04^a$	$2.19 \pm 0.21^{ab}$	$6.74 \pm 0.68^{ab}$
SA-12	$8.33 \pm 0.13^a$	$2.82 \pm 0.26^c$	$8.76 \pm 0.87^b$

The values are means ( $n = 3$ ) with standard errors.

EC electrical conductivity.

Different letters in the same column represent significant differences at  $P < 0.05$  level among sampling sites.

macro-aggregates ( $> 0.25$  mm) increased along the chronosequence of the *S. alterniflora* invasion, but decreased for fine aggregates ( $< 0.106$  mm), and showed no significant change in medium aggregates (0.106–0.25 mm) in both soil layers (Figure 2A, C). Meanwhile, soil aggregate stability (as quantified by the mean weight diameter, MWD) increased linearly with the chronosequence of the *S. alterniflora* invasion in surface layer ( $R^2 = 0.99$ ,  $P < 0.01$ ) and deep layer ( $R^2 = 0.93$ ,  $P = 0.03$ , Figure 2B, D).

### SOC and STN Concentrations in the Whole Soil and Aggregate Fractions

Along the chronosequence following *S. alterniflora* invasion, SOC and STN concentrations linearly increased by twofold in the whole soil and aggregate fractions, but the C/N ratio had no significant trend (Figure 3). The concentration of SOC in the whole soil increased from  $6.73 \pm 0.67$   $\text{g C kg}^{-1}$  soil in bare flats to  $20.31 \pm 1.03$   $\text{g C kg}^{-1}$  soil in the 12-year invaded soil by *S. alterniflora* (SA-12) in surface layer (0–20 cm) and increased from  $6.00 \pm 0.46$  to  $13.26 \pm 0.86$   $\text{g C kg}^{-1}$  soil in deep layer (20–40 cm, Figure 3C). Changes in N concentration paralleled those in C with an increase from  $0.86 \pm 0.06$  to  $2.28 \pm 0.25$   $\text{g N kg}^{-1}$  soil in surface layer and an increase from  $0.75 \pm 0.03$  to  $1.55 \pm 0.08$   $\text{g N kg}^{-1}$  soil in deep layer (Figure 3F). Within aggregate fractions, the largest SOC concentration was observed in the large macro-aggregates ( $> 1$  mm), which increased from  $10.82 \pm 1.27$  to  $24.10 \pm 1.64$   $\text{g C kg}^{-1}$  soil in surface layer and from  $10.52 \pm 0.22$  to  $16.27 \pm 0.89$   $\text{g C kg}^{-1}$  soil in deep layer. The smallest SOC concentration occurred in fine aggregates ( $< 0.106$  mm), which increased from  $5.73 \pm 0.77$  to  $15.09 \pm 0.89$   $\text{g C kg}^{-1}$  soil in surface layer and from  $4.76 \pm 0.26$  to  $10.47 \pm 1.02$   $\text{g C kg}^{-1}$  soil in

**Table 2.** Variation in  $\delta^{13}\text{C}$  Values of SOC Within Different Soil Fractions in *Spartina alterniflora* (SA) Communities and Bare Flat (BF) at Dongtan Wetlands, Shanghai, China

0–20 cm				
	BF $\delta^{13}\text{C}$ (‰)	SA-4 $\delta^{13}\text{C}$ (‰)	SA-8 $\delta^{13}\text{C}$ (‰)	SA-12 $\delta^{13}\text{C}$ (‰)
> 1 mm	– 23.99 ± 0.28aA	– 20.14 ± 0.70aB	– 19.70 ± 0.31aBC	– 18.57 ± 0.31aC
0.5–1 mm	– 23.89 ± 0.07aA	– 21.00 ± 0.56abB	– 20.32 ± 0.38aBC	– 19.06 ± 0.28abC
0.25–0.5 mm	– 23.83 ± 0.30aA	– 21.59 ± 0.50abB	– 20.41 ± 0.44aBC	– 19.36 ± 0.38bC
0.106–0.25 mm	– 24.05 ± 0.32aA	– 21.77 ± 0.56abB	– 20.37 ± 0.36aBC	– 19.15 ± 0.54abC
< 0.106 mm	– 23.82 ± 0.06aA	– 22.81 ± 0.34bA	– 21.18 ± 0.49aB	– 20.70 ± 0.41cB
Whole soil	– 23.83 ± 0.13A	– 21.94 ± 0.49B	– 20.58 ± 0.43C	– 19.38 ± 0.31D
20–40 cm				
	BF $\delta^{13}\text{C}$ (‰)	SA-4 $\delta^{13}\text{C}$ (‰)	SA-8 $\delta^{13}\text{C}$ (‰)	SA-12 $\delta^{13}\text{C}$ (‰)
> 1 mm	– 24.28 ± 0.18aA	– 22.74 ± 0.25aB	– 20.71 ± 0.52aC	– 19.44 ± 0.34aD
0.5–1 mm	– 24.22 ± 0.27aA	– 22.89 ± 0.07abB	– 21.13 ± 0.68abC	– 20.02 ± 0.33aC
0.25–0.5 mm	– 24.33 ± 0.29aA	– 23.41 ± 0.16cbA	– 21.30 ± 0.81abB	– 20.45 ± 0.52aB
0.106–0.25 mm	– 24.42 ± 0.18aA	– 23.31 ± 0.17cbA	– 20.98 ± 0.89abB	– 20.00 ± 0.55aB
< 0.106 mm	– 23.87 ± 0.04aA	– 23.66 ± 0.08cA	– 23.04 ± 0.03bB	– 21.87 ± 0.42bB
Whole soil	– 23.97 ± 0.07A	– 23.50 ± 0.13A	– 22.22 ± 0.35B	– 20.72 ± 0.33C

*The values are means with standard errors (n = 3). Different lowercase letters among soil fractions indicate significant differences at P < 0.05. Different uppercase letters indicate significant differences at P < 0.05 level among sampling sites.*

deep layer. Similar patterns were observed in STN concentration in soil fractions.

To evaluate the changes in SOC and STN in soil aggregate fractions, we calculated the mean accumulation rate of SOC and STN among different aggregate size classes. SOC accumulation rate increased along the chronosequence of *S. alterniflora* invasion for all fractions in surface soil layer, although the trends were not all statistically significant. However, in the deep soil layer, the highest SOC accumulation rate was observed in the period of 4–8 years except for fine aggregates, which increased with the invasion time. N accumulation rate showed a similar pattern as C accumulation rate (Figure S1).

The concentrations of SOC and STN in the whole soil were strongly linked to the soil aggregate stability (MWD). Soil organic C concentration increased linearly with MWD in 0–20 cm ( $R^2 = 0.78$ ,  $P < 0.01$ ) and 20–40 cm ( $R^2 = 0.80$ ,  $P < 0.01$ ) soil layer (Figure 4A). Changes in N concentration paralleled those of SOC in 0–20 cm layer ( $R^2 = 0.59$ ,  $P < 0.01$ ) and 20–40 cm layer ( $R^2 = 0.81$ ,  $P < 0.01$ , Figure 4B).

### *Spartina alterniflora*-Derived C in the Whole Soil and Aggregate Fractions

The proportion of *S. alterniflora*-derived C increased along the chronosequence following *S. alterniflora* invasion (Figure 5). Soil organic C derived from *S. alterniflora* increased from 1.60 to 9.13 g C kg<sup>-1</sup> soil in 0–20 cm layer and 0.31 to 4.31 g C kg<sup>-1</sup> soil in 20–40 cm layer after the 4–12 years of plant invasion, accounting for 18.96–40.24% and 4.66–32.04% of total SOC in the whole soil, respectively (Figures 5 and 6). The relative proportion of *S. alterniflora*-derived C was higher in 0–20 cm than 20–40 cm layer (Figure 5B, D). *S. alterniflora*-derived C in all aggregate fractions increased along the chronosequence. The largest proportion of *S. alterniflora*-derived C was observed in large macro-aggregates (> 1 mm), and the smallest was in the fine aggregates (< 0.106 mm, Figure 5A, C).

To evaluate the changes in *S. alterniflora*-derived C in soil fractions, we calculated the mean accumulation rate of *S. alterniflora*-derived C among different aggregate size classes during the three invasion




Figure 2. Proportion of water-stable aggregates along *S. alterniflora* invasion chronosequence in 0–20 cm (A) and 20–40 cm (B) soil layers, and influence of invasion year on soil aggregate stability (mean weight diameter; MWD, C and D). Vertical bars denote standard errors of means ( $n = 3$ ). Different letters over the bars indicate significant differences among sampling site at  $P < 0.05$ .

stages (0–4, 4–8, and 8–12 years). The highest accumulation rate was observed in the 8- to 12-year stage in surface soil layer, whereas the smallest occurred in the 0- to 4-year stage in the deep soil layer (Figure S2). The accumulation rate in macro- and medium aggregates showed no significant difference between the 0- to 4-year and 4- to 8-year stages, but increased at the 8- to 12-year stage in the surface layer. In the deep layer, the accumulation rate in large macro-aggregates increased significantly from  $0.39 \pm 0.03$  to  $0.91 \pm 0.08$  g C kg<sup>-1</sup> soil y<sup>-1</sup>, and the same patterns were observed in other size aggregates. However, there is no apparent difference between the 4- to 8-year and 8- to 12-year stages in macro- and medium aggregates except for the fine aggregates (Figure S2).

## DISCUSSION

### Effects of *S. alterniflora* Invasion on Soil Aggregate Stability

Soil aggregate stabilization is driven by both abiotic and biotic factors, which are related to soil properties, vegetation characteristics, land management, and climate (Bronick and Lal 2005; Cécillon and others 2010). Our results showed that soil aggregate stability increased with invasion time of *S. alterniflora* in Dongtan coastal wetlands (Figure 2). Generally, vegetation characteristics illustrated the positive effects on soil aggregate stability, which was potentially driven by above- and below-ground linkage. Above-ground stems, leaves, and litters from plants protect the soil from being ero-




Figure 3. Concentration of SOC, STN, and C/N ratio in soil fractions in 0–20 cm (A, D, G) and 20–40 cm (B, E, H) soil layers, and whole soil (C, F, I). Vertical bars denote standard errors of means ( $n = 3$ ). Different letters over the bars indicate significant differences among sampling site at  $P < 0.05$ .

ded by dissipating the wave energy (Duarte and others 2013) and counteracting the raindrop energy of rainfall (Zuazo and Pleguezuelo 2008). Meanwhile, organic compounds, which are left over from litter decomposition, could hold soil aggregates together as stabilizing agents (Liao and others 2006; Abiven and others 2009). Below-ground roots facilitate the formation and stabilization of soil aggregates through a variety of ways, including providing mechanical entanglement, changing soil density via root penetration, and acting as a binding agent by roots exudates (Jastrow and others 1998; Denef and others 2002; Bronick and Lal 2005). For example, Six and others (2004) reported that root exudations (for example, polysaccharides) acted as binding agents to hold the macro-aggregate together. Morel and others (1991) suggested that mucilage released by roots could stick soil particles together directly.

Our results also showed that macro-aggregates ( $> 0.25$  mm) increased, whereas fine aggregates ( $< 0.106$  mm) decreased along the chronose-

quence (Figure 2). This finding was consistent with the results from Zhang and others (2010), who reported that the macro-aggregates increased with *S. alterniflora* invasion in an Eastern Chinese estuarine wetland, probably attributed to greater organic matter inputs (Liao and others 2006). Macro-aggregates are generally formed around temporary binding agents such as roots, fungal hyphae, and transient agents (for example, polysaccharides, Oades and Waters 1991; Six and others 2000b). Therefore, the increasing formation of macro-aggregates in coastal wetlands could be facilitated by the increasing root biomass (Liao and others 2007; Zhang and others 2010) and soil microbial biomass (Yang and others 2016) following *S. alterniflora* invasion. The simultaneous decrease in the fine aggregate-sized fraction indicated that fine aggregates are being bound into macro-aggregates in coastal wetlands. Thus, the soil aggregate stability in coastal wetlands increased along the chronosequence of *S. alterniflora* invasion, which would be of major significance in reducing erosion, sequester-






Figure 4. SOC (A) and STN (B) versus soil aggregate stability (MWD) in 0–20 and 20–40 cm soil layers.

ing carbon, and mitigating the sea-level rise (Deegan and others 2012; Kirwan and Megonigal 2013).

### Dynamics of SOC and STN in the Whole Soil and Aggregate Fractions

The high productivity (especially roots) and structural complexity of *S. alterniflora* predispose coastal wetlands to be highly efficient in sequestering organic matters (McLeod and others 2011). Our results showed that SOC and STN concentrations in the whole soil and aggregate fractions increased along the chronosequence of *S. alterniflora* invasion, but C/N ratio had no significant changes (Figure 3). Our findings were consistent with previous studies that plant invasion profoundly influenced soil C and N pools through changes in the quantity and quality of organic matter entering soils (Windgam and Ehrenfeld 2003; Liao and others 2008). Zhang and others (2010) have reported that the above- and below-ground biomass increased with *S. alterniflora* invasion, which left distinct imprints on soil C and N pools in Eastern Chinese coastal wetlands. Organic inputs of high substrate from *S. alterniflora* induced more labile C content in the soils (Cheng and others 2008), which in turn could

enhance the recalcitrant C pool (Rovira and Vallejo 2002). Meanwhile, the residues of *S. alterniflora* were resistant to decay due to the high C/N ratio and K and Na contents (Yang and others 2009). Thus, continuous inputs, low quality of litters, and the physical protection offered by aggregates lead to SOC and STN accumulation in the whole soil and aggregate fractions along the chronosequence. Furthermore, Cheng and others (2006) documented that C and N concentrations were closely related to root distribution following *S. alterniflora* invasion. Roots and their associated microorganisms played dominant roles in soil C accumulation (Clemmensen and others 2013). However, we still lack sufficient data to differentiate the relative contribution of roots and shoots, which deserves future studies.

Our results showed that the SOC and STN concentrations in fine aggregates (< 0.106 mm) were smaller than in macro- (> 0.25 mm) and medium (0.106–0.25 mm) aggregates, which all increased with the chronosequence. This phenomenon may be induced by the process of soil aggregate formation, in which exogenous organic matter is first incorporated into the macro-aggregate fraction and the new C-enriched relative to smaller fractions (Gale and others 2000). Thus, the increasing soil aggregate stability represents an important contribution to the C and N accumulations in coastal wetlands (Figure 4). The sustained increase in SOC and STN along the chronosequence following *S. alterniflora* invasion at Dongtan wetlands suggested the large potential to sequester more C after a 12-year invasion. Therefore, the *S. alterniflora* expansion, which can sequester soil C efficiently, may be a promising strategy as an eco-engineering solution for C mitigation.

In addition to plant invasion inputs by *S. alterniflora*, we found that parts of the increased SOC were derived from extraneous inputs (Figure 6). Our finding was consistent with previous observations in a variety of coastal wetlands, indicating that the extraneous inputs mainly came from sediments or phytoplankton (Cheng and others 2006; Gebrehiwet and others 2008; Kelleway and others 2016). For example, Kelleway and others (2016) reported that sedimentation may largely control the below-ground C stocks of salt marshes, which was attributed to the allochthonous inputs and/or the increasing C preservation capacity of fine sediments. Meanwhile, Gebrehiwet and others (2008) suggested that phytoplankton partly contributed to the soil organic matter in a *S. alterniflora*-vegetated wetland. In our study, the sediments and marine phytoplankton may con-





Figure 5. Distribution proportion of SOC derived from *S. alterniflora* in soil fraction and whole soil along the chronosequence in 0–20 cm (A, B), 20–40 cm (C, D) soil layers.

tribute to the SOC accumulation to some degree, especially in the upper soil. Therefore, further studies are necessary to quantify the relative contribution of multiple sources to SOC accumulation.

### Dynamics of *S. alterniflora*-Derived C in Soil Aggregate Fractions

*Spartina alterniflora*-derived C increased along the chronosequence, but exhibited different patterns in soil aggregate fractions (Figures 5, 6). The highest proportion of *S. alterniflora*-derived C occurred in coarse macro-aggregates (> 1 mm), and the lowest was in the fine fraction (< 0.106 mm) both in 0–20 and 20–40 cm soil layers (Figures S3–S7). The decreased proportion of *S. alterniflora*-derived C from the coarse to the fine fractions indicated that

*S. alterniflora* residues were initially incorporated into the coarse fraction and then turned into the fine fraction in coastal wetlands (Desjardins and others 2006; Zhang and others 2010), achieving the long-term SOC sequestration. The fresh residues of *S. alterniflora* derived the formation of macro-aggregates, becoming coarse intra-aggregate particulate organic matter (POM). The intra-aggregate particulate POM was encrusted with microbial products and soil clay particulates, forming micro-aggregates within macro-aggregates. Previous studies have shown that C within silt and clay was more stable and highly humified (Balesdent and others 1988), and thus, the increasing rate of *S. alterniflora*-derived C in fine fractions was relatively low.



**Figure 6.** Concentration of SOC derived from *S. alterniflora* and extraneous input along the chronosequence. Different upper letters over the bars indicate significant differences of *S. alterniflora*-derived C among sampling site at  $P < 0.05$ . Different lower letters over the bars indicate significant differences of extraneous input-derived C among sampling site at  $P < 0.05$ .

The *S. alterniflora*-derived C in macro-aggregates represented an important contribution to C accumulation, even though this pool generally does not exhibit long-term stability (Tisdall and Oades 1982). Numerous studies have shown that macro-aggregates are sensitive to changes in SOM because soil structure is dependent on live binding agents (Christensen 2001; Six and others 2002; Liao and others 2006). However, macro-aggregates induced compartmentalization of substrate and soil microbes and blocked diffusion of oxygen into macro-aggregates (Killham and others 1993; Six and Jastrow 2002), thus slowing down the decomposition rate of original inputs. Meanwhile, the increasing macro-aggregates would accelerate the formation of micro-aggregates, which would be more stable than macro-aggregates (Tisdall and Oades 1982; Six and others 2000b).

The physical protection of organic matter in soil micro-aggregates was greater than that provided by macro-aggregates (Krull and others 2003). Jastrow and others (1996) found that the average turnover

time of organic C in soil micro-aggregates was 412 years, which is considerably longer than that in macro-aggregates with only 140 years. Meanwhile, the silt and clay content influenced the physical protection indirectly by affecting aggregate dynamics (Six and others 2002). Previous studies documented that physical protection of organic C within soil aggregates increased with clay content (Franzluebbers and Arshad 1997). Buyanovsky and others (1994) calculated that the mean residence time of C associated with silt and clay was more than 1000 years. The silt and clay fraction was thus regarded as a long-term pool for soil C storage (Liao and others 2006). In our study, the *S. alterniflora*-derived C accumulation rates increased along the chronosequence of plant invasion in micro-aggregates and silt + clay fractions ( $< 0.25$  mm). These results indicated that *S. alterniflora*-derived C was protected by stabilization within soil micro-aggregates and in association with silt + clay fractions. Therefore, the structural protection may play a key role in SOC sequestration in coastal wetlands.

## Implications

Our results from the chronosequence of *S. alterniflora* invasion may provide some insights into the regulation of soil aggregates on C accumulation and offer implications for conservation of coastal wetlands and development of future earth system models. In this study, soil C sequestration was largely derived from internal source inputs (that is, plants) in Dongtan coastal wetlands, and the enhanced soil C storage following *S. alterniflora* invasion was linked with improvement of soil structure. If increasing C sequestration is a strategic goal, conservation and management intervention should be prioritized for the vegetation of coastal ecosystems, as massive coastal reclamation, engineering, and urbanization dramatically degrade biodiversity and destroy coastal wetlands (Green and Short 2003; Ma and others 2014).

Our study also pointed out the importance of soil structure on SOC accumulation in coastal ecosystems, which was usually overlooked in current Earth system models. Such models (for example, Century) are built around pools that cannot be separated physically or chemically, or even quantified directly (Jenkinson and Coleman 2008; Koven and others 2013; Luo and others 2015). Recently, Abramoff and others (2018) proposed a conceptual model with five measurable soil C pools, in which aggregate C was included as well as mineral-associated organic matter, particulate organic matter, low molecular weight C, and micro-

bial biomass C. Our study highlights the suitability using aggregate C for upscaling field-based measurements to regional blue C inventories and for modeling soil C stocks across space and through time under global change scenarios.

#### ACKNOWLEDGEMENTS

This study was carried out at the Chongming Dongtan Nature Reserve, Shanghai. We acknowledge the practical help provided by Dan Wang, Qin Wang, and Haiqiang Guo. We thank Ming Li, Zhenggang Du, Yuanyuan Nie, and Xi Yang for assistance with field work. We would also like to thank anonymous reviewers for their critical comments/suggestions which improved the quality of the manuscript. This research was financially supported by the National Natural Science Foundation of China (Grant Nos. 31770559, 31370489, and 31600352), the Program for Professor of Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Learning, and “Thousand Young Talents” Program in China.

#### REFERENCES

- Abiven S, Menasseri S, Chenu C. 2009. The effects of organic inputs over time on soil aggregate stability: a literature analysis. *Soil Biol Biochem* 41:1–12.
- Abramoff R, Xu X, Hartman M, O'Brien S, Feng W, Davidson E, Finzi A, Moorhead D, Schimel J, Torn M, Mayes MA. 2018. The Millennial model: in search of measurable pools and transformations for modeling soil carbon in the new century. *Biogeochemistry* 137:51–71.
- Agrawal A, Nepstad D, Chhatre A. 2011. Reducing emissions from deforestation and forest degradation. *Ann Rev Environ Resour* 36:373–96.
- Alongi DM. 2002. Present state and future of the world's mangrove forests. *Environ Conserv* 29:33–49.
- An SQ, Gu BH, Zhou CF, Wang ZS, Deng ZF, Zhi YB, Li HL, Chen L, Yu DH, Liu YH. 2007. *Spartina* invasion in China: implications for invasive species management and future research. *Weed Res* 47:183–91.
- Balesdent J, Wagner GH, Mariotti A. 1988. Soil organic matter turnover in long-term field experiments as revealed by carbon-13 natural abundance. *Soil Sci Soc Am J* 52:118–24.
- Barto EK, Alt F, Oelmann Y, Wilcke W, Rillig MC. 2010. Contributions of biotic and abiotic factors to soil aggregation across a land use gradient. *Soil Biol Biochem* 42:2316–24.
- Bronick CJ, Lal R. 2005. Soil structure and management: a review. *Geoderma* 124:3–22.
- Bull ID, van Bergen PF, Bol R, Brown S, Gledhill AR, Gray AJ, Harkness DD, Woodbury SE, Evershed RP. 1999. Estimating the contribution of *Spartina anglica* biomass to saltmarsh sediments using compound specific carbon isotope measurements. *Org Geochem* 30:477–83.
- Buyanovsky GA, Aslam M, Wagner GH. 1994. Carbon turnover in soil physical fractions. *Soil Sci Soc Am J* 58:1167–73.
- Cécillon L, de Mello NA, De Danieli S, Brun JJ. 2010. Soil macroaggregate dynamics in a mountain spatial climate gradient. *Biochemistry* 97:31–43.
- Chambers LG, Guevara R, Boyer JN, Troxler TG, Davis SE. 2016. Effects of salinity and inundation on microbial community structure and function in a mangrove peat soil. *Wetlands* 36:361–71.
- Chen ZY, Li B, Zhong Y, Chen JK. 2004. Local competitive effects of introduced *Spartina alterniflora* on *Scirpus mariqueter* at Dongtan of Chongming Island, the Yangtze River estuary and their potential ecological consequences. *Hydrobiologia* 528:99–106.
- Cheng XL, Luo YQ, Chen JQ, Lin GH, Chen JK, Li B. 2006. Short-term  $C_4$  plant *Spartina alterniflora* invasions change the soil carbon in  $C_3$  plant dominated tidal wetlands on a growing estuarine island. *Soil Biol Biochem* 38:3380–6.
- Cheng XL, Chen JQ, Luo YQ. 2008. Assessing the effects of short-term *Spartina alterniflora* invasion on labile and recalcitrant C and N pools by means of soil fractionation and stable C and N isotopes. *Geoderma* 145:177–84.
- Chiang PN, Wang MK, Chiu CY, King HB, Hwong JL. 2004. Change in the grassland-forest boundary at Ta-Ta-Chia long term ecological research (LTER) site detected by stable isotope ratio of soil organic matter. *Chemosphere* 54:217–24.
- Chmura GL, Anisfeld SC, Cahoon DR, Lynch JC. 2003. Global carbon sequestration in tidal, saline wetland soils. *Glob Biogeochem Cycles* 17:1111.
- Choi Y, Wang Y. 2004. Dynamics of carbon sequestration in a coastal wetland using radiocarbon measurements. *Glob Biogeochem Cycles* 18:4016.
- Christensen BT. 2001. Physical fractionation of soil and structural and functional complexity in organic matter turnover. *Eur J Soil Sci* 52:345–53.
- Clemmensen KE, Bahr A, Ovaskainen O, Dahlberg A, Ekblad A, Wallander H, Stenlid J, Finlay RD, Wardle DA, Lindahl BD. 2013. Roots and associated fungi drive long-term carbon sequestration in boreal forest. *Science* 339:1615–18.
- Deegan LA, Johnson DS, Warren RS, Peterson BJ, Fleeger JW, Fagherazzi S, Wollheim WM. 2012. Coastal eutrophication as a driver of salt marsh loss. *Nature* 490:388–92.
- Denef K, Six J, Merckx R, Paustian K. 2002. Short-term effects of biological and physical forces on aggregate formation in soils with different clay mineralogy. *Plant Soil* 246:185–200.
- Desjardins T, Carneirofilho A, Mariotti A, Chauvel A, Girardin C. 1996. Changes of the forest-savanna boundary in Brazilian Amazonia during the Holocene revealed by stable isotope ratio of soil organic carbon. *Oecologia* 108:749–56.
- Desjardins T, Folgarait PJ, Pando-Bahuon A, Girardin C, Lavelle P. 2006. Soil organic matter dynamics along a rice chronosequence in north-eastern Argentina: evidence from natural  $^{13}C$  abundance and particle size fractionation. *Soil Biol Biochem* 38:2753–61.
- Donato DC, Kauffman JB, Murdiyarso D, Kurnianto S, Stidham M, Kanninen M. 2011. Mangroves among the most carbon-rich forests in the tropics. *Nat Geosci* 4:293–7.
- Duarte CM, Middelburg J, Caraco N. 2005. Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences* 2:1–8.
- Duarte CM, Losada IJ, Hendriks IE, Mazarrasa I, Marba N. 2013. The role of coastal plant communities for climate change mitigation and adaptation. *Nat Clim Change* 3:961–8.

- Duiker SW, Rhoton FE, Torrent J, Smeck NE, Lal R. 2003. Iron (hydr)oxide crystallinity effects on soil aggregation. *Soil Sci Soc Am J* 67:606–11.
- Fourqurean JW, Duarte CM, Kennedy H, Marbà N, Holmer M, Mateo MA, Apostolaki ET, Kendrick GA, Krause-Jensen D, McGlathery KJ. 2012. Seagrass ecosystems as a globally significant carbon stock. *Nat Geosci* 5:505–9.
- Franzluebbers AJ, Arshad MA. 1997. Particulate organic carbon content and potential mineralization as affected by tillage and texture. *Soil Sci Soc Am J* 61:1382–6.
- Gale W, Cambardella C, Bailey T. 2000. Root-derived carbon and the formation and stabilization of aggregates. *Soil Sci Soc Am J* 64:201–7.
- Gebrehiwet T, Koretsky MC, Krishnamurthy RV. 2008. Influence of *Spartina* and *Juncus* on salt marsh sediments. III. Organic geochemistry. *Chem Geol* 255:114–19.
- Golchin A, Baldock JA, Oades JM. 1997. A model linking organic matter decomposition, chemistry, and aggregate dynamics. In: Lal R, Kimble JM, Follett RF, Stewart BA, Eds. *Soil processes and the carbon cycle*. New York: CRC Press. p 245–66.
- Graves CJ, Makrides EJ, Schmidt VT, Giblin AE, Cardon ZG, Rand DM. 2016. Functional responses of salt marsh microbial communities to long-term nutrient enrichment. *Appl Environ Microbiol* 82:2862–71.
- Green EP, Short FT. 2003. *World atlas of seagrasses*. Berkeley (CA): California University Press.
- Harvey HR, Mannino A. 2001. The chemical composition and cycling of particulate and macromolecular dissolved organic matter in temperate estuaries as revealed by molecular organic tracers. *Org Geochem* 32:527–42.
- Hemminga MA, Cattrijsse A, Wielemaker A. 1996. Bedload and nearbed detritus transport in a tidal saltmarsh creek. *Estuar Coast Shelf Sci* 42:55–26.
- IPCC. 2013. *Climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*. Cambridge: Cambridge University Press.
- Jastrow JD. 1996. Soil aggregate formation and the accrual of particulate and mineral-associated organic matter. *Soil Biol Biochem* 28:665–76.
- Jastrow JD, Boutton TW, Miller RM. 1996. Carbon dynamics of aggregate-associated organic matter estimated by carbon-13 natural abundance. *Soil Sci Soc Am J* 60:801–7.
- Jastrow JD, Miller RM, Lussenhop J. 1998. Contributions of interacting biological mechanisms to soil aggregate stabilization in restored prairie. *Soil Biol Biochem* 30:905–16.
- Jenkinson DS, Coleman K. 2008. The turnover of organic carbon in subsoils. Part 2. Modelling carbon turnover. *Eur J Soil Sci* 59:400–13.
- Kelleway JJ, Saintilan N, Macreadie PI, Ralph PJ. 2016. Sedimentary factors are key predictors of carbon storage in SE Australian saltmarshes. *Ecosystems* 19:865–80.
- Kemper WD, Rosenau RC. 1986. Aggregate stability and size distribution. In: Klute A, Ed. *Methods of Soil Analysis. Part 1. 2nd edn Agron. Monogr. 9*, Madison (WI): ASA. p 425–42.
- Kennedy H, Beggins J, Duarte CM et al. 2010. Seagrass sediments as a global carbon sink: isotopic constraints. *Global Biogeochem Cycles* 24:57. <https://doi.org/10.1029/2010gb003848>.
- Killham K, Amato M, Ladd JN. 1993. Effect of substrate location in soil and soil pore-water regime on carbon turnover. *Soil Biol Biochem* 25:57–62.
- Kirwan ML, Megonigal JP. 2013. Tidal wetland stability in the face of human impacts and sea-level rise. *Nature* 504:53–60.
- Koven CD, Riley WJ, Subin ZM et al. 2013. The effect of vertically resolved soil biogeochemistry and alternate soil C and N models on C dynamics of CLM4. *Biogeosciences* 10:7109–31.
- Kristensen E, Bouillon S, Dittmar T, Marchand C. 2008. Organic carbon dynamics in mangrove ecosystems: a review. *Aquat Bot* 89:201–19.
- Krull ES, Baldock JA, Skjemstad JO. 2003. Importance of mechanisms and processes of the stabilisation of soil organic matter for modelling carbon turnover. *Funct Plant Biol* 30:207–22.
- Liao JD, Boutton TW, Jastrow JD. 2006. Storage and dynamics of carbon and nitrogen in soil physical fractions following woody plant invasion of grassland. *Soil Biol Biochem* 38:3184–96.
- Liao C, Luo Y, Jiang L, Zhou X, Wu X, Fang C, Chen J, Li B. 2007. Invasion of *Spartina alterniflora* enhanced ecosystem carbon and nitrogen stocks in the Yangtze estuary, China. *Ecosystems* 10:1351–61.
- Liao C, Peng R, Luo Y, Zhou X, Wu X, Fang C, Chen J, Li B. 2008. Altered ecosystem carbon and nitrogen cycles by plant invasion: a meta-analysis. *New Phytol* 177:706–14.
- Lin GH, Ehleringer JR, Rygielwicz PT, Johnson MG, Tingey DT. 1999. Elevated CO<sub>2</sub> and temperature impacts on different components of soil CO<sub>2</sub> efflux in Douglas-fir terracosms. *Glob Change Biol* 5:157–68.
- Luo Y, Ahlström A, Allison SD et al. 2015. Towards more realistic projections of soil carbon dynamics by earth system models. *Glob Biogeochem Cycles*. <https://doi.org/10.1002/2015gb005239>.
- Ma Z, Melville DS, Liu J, Chen Y, Yang H, Ren W, Zhang Z, Piersma T, Li B. 2014. Rethinking China's new great wall. *Science* 346:912–14.
- Mcleod E, Chmura GL, Bouillon S, Salm R, Björk M, Duarte C, Lovelock CE, Schlesingers WH, Silliman BR. 2011. A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO<sub>2</sub>. *Front Ecol Environ* 7:362–70.
- Morel JL, Habib L, Plantureux S. 1991. Influence of maize root mucilage on soil aggregate stability. *Plant Soil* 136:111–19.
- Neubauer SC. 2008. Contributions of mineral and organic components of tidal freshwater marsh accretion. *Estuar Coast Shelf Sci* 78:78–88.
- Oades JM, Waters AG. 1991. Aggregate hierarchy in soils. *Aust J Soil Res* 29:815–28.
- Qin P, Zhong CX. 1992. *Applied studies on Spartina*. London: Ocean Press.
- Roulet NT. 2000. Peatlands, carbon storage, greenhouse gases, and the Kyoto Protocol: prospects, and significance for Canada. *Wetlands* 20:605–15.
- Rovira P, Vallejo VR. 2002. Labile and recalcitrant pools of carbon and nitrogen in organic matter decomposing at different depths in soil: an acid hydrolysis approach. *Geoderma* 107:109–41.
- Salome C, Nunan N, Poteau R, Lerch TZ, Chenu C. 2010. Carbon dynamics in topsoil and in subsoil may be controlled by different regulatory mechanisms. *Glob Change Biol* 16:416–26.

- Six J, Jastrow JD. 2002. Organic matter turnover. In: Lal T, Ed. Encyclopedia of soil science. New York: Marcel Dekker. p 936–42.
- Six J, Elliott ET, Paustian K, Doran JW. 1998. Aggregation and soil organic matter accumulation in cultivated and native grassland soils. *Soil Sci Soc Am J* 62:1367–76.
- Six J, Paustian K, Elliott ET, Combrink C. 2000a. Soil structure and organic matter: I. Distribution of aggregate-size classes and aggregate-associated carbon. *Soil Sci Soc Am J* 64:681–9.
- Six J, Elliott ET, Paustian K. 2000b. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biol Biochem* 32:2099–103.
- Six J, Conant RT, Paul EA, Paustian K. 2002. Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant Soil* 241:155–76.
- Six J, Bossuyt H, Degryze S, Denef K. 2004. A history of research on the link between (micro) aggregates, soil biota, and soil organic matter dynamics. *Soil Tillage Res* 79:7–31.
- Sun T, Chen Q, Chen Y, Cruse R, Li X, Song C, Kravchenko Y, Zhang X. 2014. A novel soil wetting technique for measuring wet stable aggregates. *Soil Tillage Res* 141:19–24.
- Tisdall JM, Oades JM. 1982. Organic matter and water-stable aggregates in soils. *J Soil Sci* 33:141–63.
- Wang Q, An SQ, Ma ZJ, Zhao B, Chen JK, Li B. 2006. Invasive *Spartina alterniflora*: biology, ecology, and management. *Acta Phytotaxon Sin* 44:559–88.
- Wang D, Zhang R, Xiong J, Guo HQ, Zhao B. 2015. Contribution of invasive species *Spartina alterniflora* to soil organic carbon pool in coastal wetland: stable isotope approach. *Chin J Plant Ecol* 39:941–9 (in Chinese, with English abstract).
- Wilson GWT, Rice CW, Rillig MC, Springer A, Hartnett DC. 2009. Soil aggregation and carbon sequestration are tightly correlated with the abundance of arbuscular mycorrhizal fungi: results from long-term field experiments. *Ecol Lett* 12:452–61.
- Windham L, Ehrenfeld JG. 2003. Net impact of a plant invasion on nitrogen-cycling processes within a brackish tidal marsh. *Ecol Appl* 13:883–97.
- Wooller M, Smallwood B, Jacobson M, Fogel M. 2003. Carbon and nitrogen stable isotopic in *Laguncularia racemosa* (L.) (white mangrove) from Florida and Belize: implications for trophic level studies. *Hydrobiologia* 499:13–23.
- Yang SG, Li JH, Zheng Z, Meng Z. 2009. Characterization of *Spartina alterniflora* as feedstock for anaerobic digestion. *Bio-mass Bioenergy* 33:597–602.
- Yang W, Jeelani N, Leng X, Cheng XL, An XQ. 2016. *Spartina alterniflora* invasion alters soil microbial community composition and microbial respiration following invasion chronosequence in a coastal wetland of China. *Sci Rep* 6:26880. <https://doi.org/10.1038/srep26880>.
- Zhang YH, Ding WX, Luo JF, Donnison A. 2010. Changes in soil organic carbon dynamics in an Eastern Chinese coastal wetland following invasion by a C<sub>4</sub> plant *Spartina alterniflora*. *Soil Biol Biochem* 42:1712–20.
- Zhao B, Yan Y, Guo HQ, He MM, Gu YJ, Li B. 2009. Monitoring rapid vegetation succession in estuarine wetland using time series MODIS-based indicators: an application in the Yangtze River Delta area. *Ecol Indic* 9:346–56.
- Zuazo VHD, Pleguezuelo CRR. 2008. Soil-erosion and runoff prevention by plant covers: a review. *Agron Sustain Dev* 28:65–86.